

UNPUBLISHED PRELIMINARY DATA

↑ PROBABLE SYNTHESIS OF PORPHINE-LIKE SUBSTANCES
DURING CHEMICAL EVOLUTION

18p

(NASA Hunt No 4-226-62)
(NASA CR-52450)

Presented

At the Conference on The Origin of Prebiological
Systems, October 27-30, 1963 in Wakula Springs, ~~Florida~~
Fla.

FACILITY FORM 902
N65 15374
(ACCESSION NUMBER)
18
(PAGES)
CR 52450
(NASA CR OR TMX OR AD NUMBER)

(THRU)
(CODE)
06
(CATEGORY)

Conf
[2]

Personal
author:

Anton Szutka, Ph.D. [1963]
18 p refs
Department of Chemistry

GPO PRICE \$

OTS PRICE(S) \$

Hard copy (HC) \$1.00

Microfiche (MF) \$0.50


no
author

255 3006

University of Detroit U.,
Detroit, Michigan 48221

INTRODUCTION

It is already a well established theory that chemical evolution, i.e. evolution on the molecular level, has preceded the appearance of the first living cell on the earth. Current concepts suggest that a relatively simple primitive atmosphere, consisting of methane, ammonia, water, and also probably some molecular hydrogen under influence of ultra-violet radiation, electrical discharges and high energy radiations has been transformed into more complicated compounds (1, 2, 3). It has been generally accepted that the temperature at the time of this transformation was less than 100°C. The raw material thus produced, accumulated in pools of water that condensed, forming an organic milieu, i.e. the non-living mixture of carbon compounds which is generally considered as a prerequisite of the origin of life. Some estimates made by Urey (2) indicate that the primitive oceans might have contained as much as 10 per cent of dissolved organic compounds. The synthesis of these raw materials continued until a state of equilibrium was reached, in which the action of available energy sources produced as much of the raw material as it destroyed. In this period of time only relatively simple molecules could be formed because the more complicated and labile ones would be destroyed by the energetic ultraviolet radiation reaching the surface of the earth, unscreened by the ozone layer, which now protects the earth.



The above theories of Oparin (1) and Urey (2) have been subjected to rigid laboratory tests by Miller (4, 5), Pavlovskaya and Pasynskii (6), Abelson (7), Oro (8), Hasselstrom (9), Paschke (10), and others (11, 12). This laboratory work has produced various products of biochemical importance, such as amino acids, aliphatic acids, polyhydroxy compounds, aldehydes and urea. Subjecting these simple molecules to further action of ultraviolet radiation or mild heating, produced a variety of complex organic molecules. Such higher order reactions, for example, result in production of polypeptides from amino acids, as demonstrated by Fox (13, 14), Akabori (15), Doty (16), and others (17), in actual laboratory experiments.

In the reviewed literature, most of the authors (1, 18, 19, 20) dealing with the problem of the origin of life, are convinced that at some stage of chemical evolution, porphyrins must have developed. However, until recently there was no laboratory evidence that porphyrins or porphine-like substances could be synthesized from precursors which were available during the period of chemical evolution on earth.

This paper presents observations having direct bearing on the formation of the organic millieu, namely the synthesis of porphine-like substances from simple precursors.

EXPERIMENTAL PROCEDURES

It has been known that pyrrole and benzaldehyde when heated

to about 180° C. condense to form $\alpha, \beta, \gamma, \delta$ -tetraphenylporphine (21) (Fig. 1) and $\alpha, \beta, \gamma, \delta$ -tetraphenylchlorine (22). The same system, with 5 ml. of pyrrole, 10 ml. of benzaldehyde, 10 ml. of pyridine (all freshly redistilled), and 5 g. of zinc acetate, was used in our first experiments (23). The pyridine serves as a solvent. Zinc acetate was found to increase the yield (22, 24). The mixture was placed in a screw-cap Pyrex tube with approximately 2 ml. of air space above the liquid and irradiated with Co^{60} γ -rays for 10 hrs. at the rate of approximately 0.5 megarepents per hour. The solution removed from the irradiation vessel, reddish in color, was diluted with chloroform, treated with an equal volume of 6 N hydrochloric acid, and stirred vigorously for 3 hours. The acidic solution was then neutralized with ammonia. The chloroform solution, separated from the aqueous layer, was washed several times with distilled water and chromatographed on activated alumina, as suggested by Priesthoff and Banks (24), with fresh chloroform as the eluent. Three chromatographic separations removed most of the tarry products from the chloroform solution, which was subsequently subjected to chromatography on a Magnesol column. The dark-green band was repacked into a second column, wetted with ether, and eluted with chloroform. The procedure was repeated until a single, clear band was obtained on the column.

In recent experiments (25), a system, consisting of 3 ml. of freshly distilled pyrrole, 6 ml. of benzaldehyde, and 4 ml.

of water, was placed in a 250 ml. beaker, and then irradiated, at a distance of 25 cm with ultraviolet light, using a 100 Watt Hanovia Utility Model Lamp. The radiation transmitted through the filter was 45 per cent at 2500 Å, 90 per cent at 3000 Å and the output of the lamp was 1.5×10^{15} quanta per sec per cm^2 or 5×10^{16} quanta per sec per area of the mixture exposed toward radiation. The doses were measured by chemical actinometry, using an uranyl oxalate actinometer according to Forbes and Heidt (26). After exposure, the reddish-brown solution was diluted with approximately 20 ml. of chloroform and washed with 20 ml. of distilled water. After separation in a separatory funnel, the aqueous layer was discarded, and the chloroform layer was transferred into a beaker charged with approximately 20 g of anhydrous sodium sulfate to remove the excess of water. The sodium sulfate was then filtered off and discarded, while the filtrate was diluted up to 50 ml. volume with chloroform and stored for the quantitative determination of porphines.

The final product separation was accomplished by column chromatography, using activated alumina (F-20) and Florosil (60-100 mesh) as adsorbents and fresh chloroform as eluent. Two chromatographic treatments on alumina and three to four chromatographic separations on Florosil yielded a pure product. The porphines appeared in the eluents and collections of eluent were continued until no Soret band appeared in the effluent.

The same mixture of pyrrole, benzaldehyde and water when placed in the dark without irradiation, also produced porphines.

The isolation and quantitative determination of porphines was done by the same methods used for irradiated mixtures.

If, instead of water, pyridine was used as a solvent in the system, and the mixture was then irradiated for two hours, no porphine-like substances could be detected by previously used methods. Aging of the mixture for ten days produced only traces of porphines.

A typical spectrum of a compound isolated from an irradiated mixture and the reference spectrum of the compound synthesized by the method of Ball, et. al. (22), are presented in Fig. 2. Both spectra are identical with those obtained by Thomas et al. (27), and Dorrough et al. (28). For quantitative estimation of the yield, the molar extinction coefficient of 18.7×10^3 at 515 m μ maximum was used, as suggested by Thomas and Martell (27). Identity of the compound was further established by the formation of a zinc chelate, according to the procedure of Rothemund and Menotti (29), and Dorrough et al. (28). Both spectra, the reference zinc chelate and the chelate of the compound isolated from the irradiated mixture are identical, as shown in Fig. 3.

It is also well known that acid salts of $\alpha, \beta, \gamma, \delta$ -tetraphenylporphines exhibit a characteristic spectra (28, 30). Therefore, part of the benzene solution of the reference compound and part of the benzene solution of compounds isolated from irradiated mixtures were evaporated to dryness and treated with glacial acetic acid. Upon recording the spectra (Fig. 4)

they were found to be identical with those listed in the literature (28, 30).

Although the amount of porphines isolated after γ -irradiation was in the microgram range, this quantity was sufficient to make positive identification.

Fig. 5 and 6 represent the total yields of porphines isolated from the stored mixtures, plotted against time of storage. Each line in these graphs represents different irradiation conditions.

DISCUSSION

In our present study we have shown that porphine-like structures can be synthesized in the presence of oxygen from precursors which were available in the very earliest stage of chemical evolution. Aldehydes were found in products of the action of electrical discharges on the Urey atmosphere in Miller's experiment (4, 5). Pyrroles and pyrrolidines were easily formed from ammonia, acetylene, and other unsaturated hydrocarbons by simple catalysis or under influence of ultra-violet radiation (1). In addition, Lichtin (31) found pyrroles as the major product of the reaction of active nitrogen with 1,3-butadienes.

The results of this study also indicate that the presence of water, or rather the suspension of the organic matter in water, increases the yield of porphine like substances considerably. Oparin (1), and Haldane (3), have already indicated

the possibility of evolution of living matter from the pools rich in organic matter, rather than on solid surfaces. The concentration of organic matter could be quite high, if part of the water solvent evaporated, as the case might be in lakes and lagoons.

The increase in the yield of porphine-like substances, on standing, shows that a process of autocatalysis takes place. This process, postulated by Calvin (19), now can be supported by experimental evidence.

A closer examination of Fig. 5 and 6 reveals that there is a difference in the total yield of porphines after standing when the mixture is originally irradiated with different kinds of electromagnetic radiation. Initial exposure to the tungsten lamp and no initial exposure both on standing in semidarkness seem to have a similar effect. The rate of increase in the production of porphines is the same within the experimental error. On the other hand, an initial irradiation with ultraviolet light has a lower rate of increase of porphine production under identical conditions of storage. This comparison has been made on aqueous suspensions of the starting materials and in the presence of oxygen.

There is a general agreement among many authors that synthesis of porphyrins is a necessary step for the origin of life. However, the question of the time of the appearance of porphyrins is subject to discussion. Gaffron (18), and Calvin (19), support the ideas that porphyrins were already active in

the very earliest quasi-living organic structures. Miller and Urey (5), consider this not a necessity and suggest that porphyrins may have arisen during the evolution of primitive organisms. Strughold and Ritter (32), consider pre-existing stores of oxygen, produced by photochemical dissociation of water, as a prerequisite for the formation and development of chlorophyll, while the heme types of porphyrins were formed about 1 to 1.5 billion years later.

Although at the present time it seems to be impossible to determine with absolute confidence at what stage of chemical evolution porphine-like substances developed, certain deductions can be based on the experimental evidence now available. It can be argued that after the stores of simpler organic compounds reached their saturation point, in the first stage of chemical evolution, there was need for a reduction of the energy level of the source in order to continue chemical evolution to a more complicated nature. Therefore the photolysis of water and the formation of an ozone layer cut out short ultraviolet radiation completely, leaving only a mild energy source of longer ultraviolet and visible radiation. Considering the lower rate of production of porphine-like substances on standing which were irradiated with ultraviolet light, it might be deduced that the formation of these substances occurred in an oxidative atmosphere.

In addition evolutionary selection pressure was exerted for the synthesis of porphine-like substances as soon as the

reductive atmosphere was converted into an oxidative one. Since with the cut off of the short ultraviolet radiation not enough energy could be provided by the long wave ultraviolet and visible radiation for certain chemical synthesis, the appearance of porphines and especially their metal chelates could be considered a necessity. Their aid in chemical transformations such as hydrogen transfer or oxidation is well established.

Also simultaneously with the change of reductive atmosphere to an oxidative one due to the photolysis of water, hydrogen peroxide appeared. If it remained in contact with organic substances, the results would be fatal. Therefore, in order to continue chemical evolution, it was necessary that a mechanism be developed for the destruction of these large amounts of hydrogen peroxide. Calvin (5) has already pointed out that incorporation of ferric iron into a heme- type molecule would increase the catalytic activity of iron for the destruction of hydrogen peroxide by a factor of 10^3 , while the addition of certain protein arrangement increases this destruction rate by an additional factor of 10^5 .

These last two arguments indicate that with the transition of a reductive atmosphere into an oxidative one, an immediate need was created for the formation of the porphine-like substances.

This work is supported in part by the Grant NsG-226-62 from the National Aeronautics and Space Administration.

REFERENCES

1. (a) A.I. Oparin, The Origin of Life on the Earth (translated by A. Synge), 3rd ed., Academic Press, New York, 1957; (b) Life, its Nature, Origin and Development (translated by A. Synge), Academic Press, New York, 1962.
2. H.C. Urey, The Planets, Yale University Press, New Haven, Connecticut, 1952.
3. J.B.S. Haldane, Collection of Essays on the Origin of Life, Penguin Books, Ltd., London, 1954.
4. (a) S. L. Miller, A production of amino acids under possible primitive earth conditions. *Science* 117, 528-529 (1953); (b) Production of some organic compounds under possible primitive earth conditions. *J. Am. Chem. Soc.* 77, 2351-2361 (1955); (c) The formation of organic compounds on the primitive earth. *Ann. N.Y. Acad. Sci.* 69, 260-274 (1957); (d) The mechanism of synthesis of amino acids by electric discharges. *Biochim. et Biophys. Acta* 23, 480-489 (1957).
5. S.L. Miller and H.C. Urey, Organic Compound Synthesis on the Primitive Earth. *Science* 130, 245-251 (1959).
6. T.E. Pavlovskaya and A.G. Pasyanskii, The original formation of amino acids under action of ultraviolet rays and electric discharges. In Proceedings of the First International Symposium on the Origin of Life on the Earth.

- A. I. Oparin, et al. ed. pp. 151-157, Pergamon Press, London, (1959).
7. P.H. Abelson, Amino acids formed in "primitive atmosphere". Science 124, 935 (1956)
 8. (a) J. Oro, Mechanism of synthesis of adenine from hydrogen cyanide under possible primitive earth conditions. Nature 191, 1193-1194 (1961); (b) J. Oro and C. L. Guidry, Direct Synthesis of polypeptides. I. Polycondensation of glycine in aqueous ammonia. Arch. Biochem. Biophys. 93, 166-171 (1961); (c) J. Oro and S.S. Kamat, Amino acid synthesis from hydrogen cyanide under possible primitive earth conditions. Nature 190, 442-443 (1961); (d) J. Oro and A.P. Kimball, Synthesis of purines under possible primitive earth conditions. I. Adenine from hydrogen cyanide. Arch. Biochem. Biophys., 94, 217-227 (1961).
 9. (a) T. Hasselstrom and M.C. Henry, New Synthesis of Oxalic Acid. Science, 123, 1038-1039 (1956); (b) T. Hasselstrom, M.C. Henry, and B. Murr, Synthesis of Amino Acids by Beta Radiation. Science, 125, 350-351 (1957)
 10. R. Paschke, R.W.H. Chang, and D. Young, Probable role of gamma irradiation on origin of life. Science 125, 881 (1957).
 11. (a) K. Bahadur, Photosynthesis of amino acids from paraformaldehyde and potassium nitrate. Nature 173, 1141 (1954); (b) K. Bahadur and R. R. Srivastava, Effect

of catalysts and the period of irradiation on the photosynthesis of amino acids in the mixture of para-formaldehyde and potassium nitrate. Zh. Obshchei Khimii 31, 3017-3020 (1961)

12. A. Palm and M. Calvin, Primordial Organic Chemistry. I. Compounds resulting from electron irradiation of $C^{14}H_4$. J. Am. Chem. Soc., 84, 2115 (1962).
13. (a) S. W. Fox, Evolution of protein molecules and thermal synthesis of biochemical substances. Am. Scientist 44, 347-362 (1956); (b) S.W. Fox, How did life begin? Science 132, 200-208 (1960).
14. S.W. Fox and K. Harada, Thermal copolymerization of amino acids in the presence of phosphoric acids. Arch. Biochem. Biophys. 86, 281-285 (1960).
15. S. Akabori, On the origin of the fore-protein. In Proc. of the First International Symposium on the Origin of Life on the Earth (A.I. Oparin et al., eds. pp. 189-196, Pergamon Press, London, (1959).
16. P. Doty, K. Imahori, and E. Klemperer, The solution, properties and configurations of a polyampholytic polypeptide, copoly - L - lysine - L - glutamic acid. Proc. Nat. Acad. Sci. 44, 424-431 (1958).
17. (a) G. Ferrari and R. Cultrera, Sul meccanismo della sintesi fotochimica di amino acidi. Nota I. Gazzetta Chim. Ital. 90, 1522-1529 (1960); (b) G. Ferrari and R. Cultrera, Sul meccanismo nos della sintesi fotochimica di amino

- acidi. Nota II. Gazz. Chim. Ital. 90, 1637-1644 (1960)
18. H. Gaffron, The origin of life. Perspectives in Biol. Med. 3, 163-212 (1960)
 19. (a) M. Calvin, Evolution of enzymes and the photosynthetic apparatus. Science 139, 1170-1174 (1959); (b) M. Calvin, The Chemistry of Life. Chem. Eng. News 39, 96-104 (May 22, 1961); (c) M. Calvin, The origin of life on earth and elsewhere. Perspectives in Biol. Med. 5, 399-422 (1962); (d) M. Calvin, Evolution of photosynthetic mechanisms, Perspectives in Biol. Med. 5, 147-172 (1962).
 20. C. Sagan, On the origin and planetary distribution of life. Radiation Res. 15, 174-192 (1961)
 21. P. Rothemund, New porphyrin synthesis. The synthesis of porphine. J. Am. Chem. Soc. 58, 625-627 (1936).
 22. R. H. Ball, G.D. Dorough, and M. Calvin, A further study of the porphine-like products of the reaction of benzaldehyde and pyrrole. J. Am. Chem. Soc. 68, 2278-2281 (1946).
 23. A. Szutka, J. F. Hazel and W.M. McNabb, Synthesis of Porphine-like substances induced by gamma-irradiation. Radiation Res. 10, 597-603 (1959)
 24. J. H. Priesthoff and C.V. Bauks, A new method of purifying $\alpha, \beta, \gamma, \delta$ -tetraphenylporphines. J. Am. Chem. Soc. 76, 937-938 (1954).
 25. A. Szutka, Porphine-like substances: Probable synthesis during chemical evolution. Science (submitted for publication).

26. G.S. Forbes and L. J. Heidt, Optimum composition of uranyl oxalate solutions for actimometry. J. Am. Chem. Soc. 56, 2363 (1934).
27. D. W. Thomas and A. E. Martell, Absorption spectra of para-substituted tetraphenylporphines. J. Am. Chem. Soc. 78, 1338-1343 (1956).
28. G.D. Dorough, J.R. Miller, and F.M. Huennekens, Spectra of the metallo-derivatives of $\alpha, \beta, \gamma, \delta$ -tetraphenylporphine. J. Am. Chem. Soc. 73, 4315-4320 (1951).
29. P. Rothmund and A.M. Menotti, Porphyrin studies. V. The metal complex salts of $\alpha, \beta, \gamma, \delta$ -tetraphenylporphine. J. Am. Chem. Soc. 70, 1808-1812 (1948).
30. C. V. Banks and R.E. Bisque, Spectrophotometric determination of zinc and other metals with $\alpha, \beta, \gamma, \delta$ -tetraphenylporphine. Anal. Chem. 29, 522-526 (1957).
31. (a) A. Tsukamoto and N.N. Lichtin, Reactions of Active nitrogen with organic substrates. I. The monomeric products of the reaction with 1, 3 - butadiene. J. Am. Chem. Soc. 84, 1601 (1962); (b) A. Tsukamoto and N.N. Lichtin, Non-degradative reaction of "active" nitrogen with conjugated dienes. J. Am. Chem. Soc. 82, 3798 (1960).
32. H. Strughold and C. L. Ritter, Oxygen production during the evolution of the earth's atmosphere. Aerospace Med. 33, 275-278 (1962).

